

SUPPLEMENTARY MATERIAL



Exploratory Association Between Recorded Study-Specific ADL and Perceived Quality-of-Life Scores Among Older Adults in One Indonesian Residential Care Facility: A Cross-Sectional Secondary Analysis

Akhmad Yanuar Fahmi Pamungkas¹, Silfin Nabila², Fany Anitarini¹

¹ Faculty of Nursing, Universitas Dr. Soekardjo, Banyuwangi, East Java, Indonesia

² Bachelor of Nursing Programme, Universitas Dr. Soekardjo, Banyuwangi, East Java, Indonesia

Correspondence: Akhmad Yanuar Fahmi Pamungkas (yanuarfahmi20@gmail.com)

Received: 16 July 2026 | **Revised:** 19 July 2026 | **Accepted:** 19 July 2026

Keywords: Activities of daily living; Exploratory analysis; Measurement uncertainty; Older adults; Perceived quality of life; Residential care

TABLE OF CONTENT

FIGURE S1 |

Operational Quality-Of-Life Categories Across Functional Categories. Categories Are Descriptive and Not Externally Validated. The Pattern Is Non-Monotonic in the Two Smallest Dependence Groups. A1_P2

FIGURE S2 |

Marginal Adjusted Perceived Quality-Of-Life Estimates Across the Observed Corrected ADL Range. Predictions Are from the Age- And Sex-Adjusted HC3 Model and Are Included as A Sensitivity Visualization Rather Than A Causal Dose-Response Curve. A1_P2




FIGURE S1 | Operational Quality-Of-Life Categories Across Functional Categories. Categories Are Descriptive and Not Externally Validated. The Pattern Is Non-Monotonic in the Two Smallest Dependence Groups.




FIGURE S2 | Marginal Adjusted Perceived Quality-Of-Life Estimates Across the Observed Corrected ADL Range. Predictions Are from the Age- And Sex-Adjusted HC3 Model and Are Included as A Sensitivity Visualization Rather Than A Causal Dose-Response Curve.